

Slide 1

Partial Pressure

When moles are not moles and
atmospheres are not atmospheres

Slide 2

Chemists are Pragmatists

Most gas phase reactions occur in sealed flasks –
you've got to keep the reactants from
escaping!

That means that, typically, the volume is fixed
and the temperature is known (unless it is a
very exothermic or endothermic reaction and
the temperature isn't controlled).

Slide 3

The Gas Law revisited

$PV = nRT$

In a sealed flask, V is constant and (usually) so is
Temperature. Collect the constants!

$P = n \frac{RT}{V} = n * \text{constant}$

The pressure is directly proportional to n, the number of
moles!

Slide 4

Suddenly bizarre units...

This is critically important for a reaction because, as you know, reactions are all about MOLES! MOLES! MOLES!

For gas phase reactions, the pressure is sometimes a substitute for the number of moles.

YOU CAN ACTUALLY MEASURE THE AMOUNT OF SOMETHING IN ATMOSPHERES!!!!!!

Slide 5

PV=nRT does not discriminate

While it may not be immediately obvious, there's another interesting thing about PV=nRT...

None of the variables directly depends on the identity of the gas molecules!

P, V, and T are physical properties of the system! Even n is just the number of particles - any particles!

Slide 6

What's it mean?

It means that the gas laws are additive!!!!

If I have a mixture of gases, I can look at the physical properties (P, V, T or n) as belonging to one gas separate from the others, or to all the gases collectively.

(Well, except for T, since the gases must all be at the same temperature.)

Slide 7

Consider Pressure:

What's pressure?

The combined effect of moving gas molecules bouncing off of things.

If you have Hydrogen and Helium mixed in a flask, the total pressure comes from the combined collisions of the Hydrogen and the Helium.

Slide 8

Keep 'em apart

Hydrogen and Helium independently obey the ideal gas equation.

$$P_{\text{He}} V = n_{\text{He}} R T$$
$$P_{\text{H}_2} V = n_{\text{H}_2} R T$$

Slide 9

Put 'em together

Hydrogen and Helium collectively obey the ideal gas equation.

$$P_{\text{total}} V = n_{\text{total}} R T$$
$$P_{\text{total}} V = (n_{\text{He}} + n_{\text{H}_2}) R T$$

Slide 10

Lump the constants..

$$P_{\text{total}} = \frac{n_{\text{total}}RT}{V}$$
$$P_{\text{total}} = \frac{(n_{\text{He}} + n_{\text{H}_2})RT}{V} = \frac{n_{\text{He}}RT}{V} + \frac{n_{\text{H}_2}RT}{V}$$
$$P_{\text{total}} = P_{\text{He}} + P_{\text{H}_2}$$

The total pressure is the sum of the **partial pressure**

Slide 11

Partial Pressure

The partial pressure is defined as the pressure exerted by a gas ignoring the presence of any other gases.

(You could also define a partial volume similarly.)

Considering that for an ideal gas, gas molecules don't interact (one of the 2 conditions), this would seem to be a logical result.

Slide 12

Dalton's Law of Partial Pressures

$$P_{\text{total}} = P_{\text{He}} + P_{\text{H}_2}$$

This is just one specific example of the more general rule:

$$P_{\text{total}} = P_1 + P_2 + P_3 + \dots$$

Where P_i is the partial pressure of gas i .

Slide 13

Calculating partial pressure

As we've seen, the partial pressure is just like any old pressure:

$$PV = nRT$$
$$P = \frac{nRT}{V}$$

Remember, we are in one flask at one temperature, so RT/V is constant

Slide 14

Pick a gas, any gas

$$P_{\text{He}} = n_{\text{He}} \frac{RT}{V}$$
$$P_{\text{total}} = n_{\text{total}} \frac{RT}{V}$$

In a flask of constant T, T and V are constant for each gas and for the combination of all gases!

Slide 15

Pick a gas, any gas

$$P_{\text{He}} = n_{\text{He}} \frac{RT}{V}$$
$$P_{\text{total}} = n_{\text{total}} \frac{RT}{V}$$
$$\frac{P_{\text{total}}}{n_{\text{total}}} = \frac{RT}{V}$$

So: $P_{\text{He}} = \frac{n_{\text{He}} P_{\text{total}}}{n_{\text{total}}} = X_{\text{He}} P_{\text{total}}$

Slide 16

Mole Fraction

χ_{He} is called the "mole fraction of He".

χ is a unit of concentration!

Slide 17

Little Bitty Problem

If I had 0.25 mol H₂ and 0.75 mol He in a 1 L flask at 273 K, what is the partial pressure of each gas and the total pressure in the flask?

$$P_{\text{H}_2} = \frac{n_{\text{H}_2} R T}{V} = \frac{0.25 \text{ mol} * 0.082056 \text{ L atm} * 273 \text{ K}}{1 \text{ L mol K}}$$
$$P_{\text{H}_2} = 5.6 \text{ atm}$$
$$P_{\text{He}} = \frac{n_{\text{He}} R T}{V} = \frac{0.75 \text{ mol} * 0.082056 \text{ L atm} * 273 \text{ K}}{1 \text{ L mol K}}$$
$$P_{\text{He}} = 16.8 \text{ atm}$$

Slide 18

$$P_{\text{tot}} = \frac{n_{\text{tot}} R T}{V} = \frac{1.0 \text{ mol} * 0.082056 \text{ L atm} * 273 \text{ K}}{1 \text{ L mol K}}$$
$$P_{\text{tot}} = 22.4 \text{ atm} = 5.6 \text{ atm} + 16.8 \text{ atm}$$

Notice, I get the same results if I start from the total pressure and divvy it up:

$$P_{\text{H}_2} = \chi_{\text{H}_2} P_{\text{tot}} = \frac{0.25 \text{ mol}}{1.0 \text{ mol}} * 22.4 \text{ atm} = 5.6 \text{ atm}$$
$$P_{\text{He}} = \chi_{\text{He}} P_{\text{tot}} = \frac{0.75 \text{ mol}}{1.0 \text{ mol}} * 22.4 \text{ atm} = 16.8 \text{ atm}$$

Slide 19

Tro Problem 5.48

A 1.0 L container of liquid nitrogen is kept in a closet measuring 1.0 m by 1.0 m by 2.0 m. Assuming that the container is completely full, that the temperature is 25.0 C, and that the atmospheric pressure is 1.0 atm, calculate the percent (by volume) of air that would be displaced if all of the liquid nitrogen evaporated. [Liquid nitrogen has a density of 0.807 g/mL]

Slide 20

Volume displaced

If you release a second gas into a room, there are two options: either the pressure increases (more moles!) or some of the air in the room escapes.

Unless the room is airtight and sealed, some of the air will escape because the pressure in the room wants to be the same as the pressure outside.

Slide 21

Volume of air

The air fills the closet.

$$V_{\text{closet}} = l \times w \times h = 1.0 \text{ m} \times 1.0 \text{ m} \times 2.0 \text{ m} \\ = 2.0 \text{ m}^3$$

Is this a good unit?

It's the BEST UNIT!!! It's the SI unit!! (But we'd probably rather have L ☺)

Slide 22

Volume of air

The air fills the closet.

$$V_{\text{closet}} = 2.0 \text{ m}^3 \cdot \frac{(100 \text{ cm})^3}{(1 \text{ m})^3} \cdot \frac{1 \text{ mL}}{1 \text{ cm}^3} \cdot \frac{1 \text{ L}}{1000 \text{ mL}}$$
$$V_{\text{closet}} = 2000 \text{ L}$$

Slide 23

How much Nitrogen?

I'm sure many of you are tempted to say 1.0 L.

1.0 L is the volume of LIQUID nitrogen. (Which means there's really only 1999 L of air in the room!)

We need to know how much GASEOUS nitrogen is formed when the liquid evaporates.

Slide 24

1.0 L of liquid is ??? Gas?

????

$PV = nRT$

What do we know?

We know the R, T, P for sure.
We want to know V
So we need to know n.

Slide 25

Finding n

We know the volume of the liquid and its density, so we know...

...the mass of the liquid which is the same as...

...the mass of the gas, which can be used to find...

...the moles of gas, by using...

...the molar mass!

Slide 26

$1.0 \text{ L N}_2 \text{ liq} \times \frac{1000 \text{ mL}}{1 \text{ L}} \times \frac{0.807 \text{ g}}{1 \text{ mL}} = 807 \text{ g N}_2 \text{ liquid}$

$807 \text{ g N}_2 \text{ liquid} = 807 \text{ g N}_2 \text{ gas}$

$807 \text{ g N}_2 \text{ gas} \times \frac{1 \text{ mol N}_2}{28.014 \text{ g N}_2} = 28.81 \text{ mol N}_2$

Slide 27

Ideal Gas Law

$PV = nRT$

$V = \frac{nRT}{P} = \frac{(28.81 \text{ mol}) (0.082058) * 298 \text{ K}}{1 \text{ atm}}$

$V = 704.5 \text{ L N}_2$

704.5 L N₂ should displace 704.5 L air

Slide 28

% Displaced

704.5 L displaced air * 100 = 35.24%
1999 L air in closet

Slide 29

Let's actually do some chemistry!

Consider a previously evacuated 1 L sealed flask at 450 C which contains 10 g of H₂ and 10 g of O₂. If I make steam, what is the final pressure in the flask?

Where do we start?

Of course, chemistry ALWAYS starts with a balanced equation!

Slide 30

Let's actually do some chemistry!

Consider a 1 L sealed flask at 450 C which contains 10 g of H₂ and 10 g of O₂. If I make steam, what is the final pressure in the flask?

$2\text{H}_2(g) + \text{O}_2(g) \rightarrow 2\text{H}_2\text{O}(g)$

And now....???

Slide 31

YES! It IS a limiting reactant problem!

$$10 \text{ g H}_2 \cdot \frac{1 \text{ mol H}_2}{2.016 \text{ g H}_2} = 4.96 \text{ mol H}_2$$
$$4.96 \text{ mol H}_2 \cdot \frac{2 \text{ mol H}_2\text{O}}{2 \text{ mol H}_2} = 4.96 \text{ mol H}_2\text{O}$$
$$10 \text{ g O}_2 \cdot \frac{1 \text{ mol O}_2}{32 \text{ g O}_2} = 0.3125 \text{ mol O}_2$$
$$0.3125 \text{ mol O}_2 \cdot \frac{2 \text{ mol H}_2\text{O}}{1 \text{ mol O}_2} = 0.625 \text{ mol H}_2\text{O}$$

Oxygen is the limiting reagent – and we make 0.625 mol H₂O.

Slide 32

And so...???

You might be ready to PV=nRT, but...

$$2\text{H}_2(g) + \text{O}_2(g) \rightarrow 2\text{H}_2\text{O}(g)$$

Note that both the reactants and products are gases. That means all 3 species will contribute to the total pressure at the end, so we need to keep track of the amounts of all of them.

Slide 33

A little molar accounting

$$2\text{H}_2(g) + \text{O}_2(g) \rightarrow 2\text{H}_2\text{O}(g)$$

I	4.96 mol	0.3125 mol	0 mol
C	-2x mol	-x mol	+2x mol
E	(4.96 - 2x)	0 (LR)	0.625 mol

x = 0.3125 mol

Slide 34

A little molar accounting

2H_2	(g)	+	O_2	(g)		$2\text{H}_2\text{O}$	(g)
I	4.96 mol		0.3125 mol			0 mol	
C	-2(0.3125)		-0.3125			+2(0.3125)	
E	4.335		0 (LR)			0.625 mol	

So the total moles of gas are:
 $4.335 + 0.625 = 4.96 \text{ mol}$

Slide 35

$PV = nRT$
 $T = 450 \text{ C} + 273.15 = 723.15 \text{ K}$

$P = \frac{n_{\text{gas}}RT}{V} = \frac{4.96 \text{ mol} * 0.082056 \text{ L atm} * 723.15 \text{ K}}{\text{mol K} \cdot 1 \text{ L}}$

$P = 294 \text{ atm}$

Slide 36

Another Chemistry problem

32.0 g of CaCO_3 (limestone) is placed in a 2.0 L previously evacuated sealed flask with 1.5 atm HCl (g) at 125 C. What is the final pressure in the flask if the following reaction is known to occur:

$\text{CaCO}_3 \text{ (s)} + 2 \text{HCl (g)} \rightarrow \text{CaCl}_2 \text{ (s)} + \text{H}_2\text{O (g)} + \text{CO}_2 \text{ (g)}$

Slide 37

Another limiting reagent problem!

$$32.0 \text{ g CaCO}_3 \times \frac{1 \text{ mol CaCO}_3}{100.09 \text{ g}} \times \frac{1 \text{ mol CO}_2}{1 \text{ mol CaCO}_3} = 0.32 \text{ mol CO}_2$$

PV=nRT for HCl

$$n = \frac{PV}{RT} = \frac{(1.5 \text{ atm})(2.0 \text{ L})}{0.08206 \text{ Latm/mol K} (125 \text{ C} + 273.15)}$$

$$n = 0.0918 \text{ mol HCl} \times \frac{1 \text{ mol CO}_2}{2 \text{ mol HCl}} = 0.0459 \text{ mol CO}_2$$

So, HCl is the limiting reagent!

Slide 38

And again with the molar accounting

$$\text{CaCO}_3 (s) + 2 \text{HCl} (g) \rightarrow \text{CaCl}_2 (s) + \text{H}_2\text{O} (g) + \text{CO}_2 (g)$$

I NA	.0918 mol	NA	0	0
C NA	-2X	NA	+x	+x
E NA	0 (LR)	NA	0.0459 mol	0.0459 mol

Slide 39

Final Pressure

$$N_{\text{total}} = n_{\text{H}_2\text{O}} + n_{\text{CO}_2} = 0.0459 \text{ mol} + 0.0459 \text{ mol} = 0.0918 \text{ mol}$$

$$P = \frac{nRT}{V} = \frac{0.0918 \text{ mol} \times 0.08206 \text{ Latm/mol K} \times 398.15 \text{ K}}{2.0 \text{ L}}$$

P = 1.5 atm

Slide 40

Clicker Question #1

2.0 g of hydrogen and 10.0 g of oxygen are added to a sealed, evacuated 10.0 L flask. The mixture is reacted at 550 °C for 6 hours. Assuming the reaction goes to completion, what is the total pressure in the flask after 6 hours (before it cools down)?

A. 6.8 atm
 B. 4.3 atm
 C. 2.1 atm
 D. 3.8 atm

Slide 41

2 g H₂ * 1 mol/2 g = 1.0 mol H₂ * (2 H₂O/2 H₂) = 1 mol H₂O
 10 g O₂ * 1 mol/32 g = 0.3125 mol O₂ * (2 mol H₂O/1 mol O₂) = 0.625 mol H₂O

2H_{2(g)} + O_{2(g)} → 2 H₂O_(g)

I 1.0 mol	0.3125 mol	0 mol
C -2x	-x	+2x
E 1-2x	0 LR	2x mol

0.3125-x = 0
 X = 0.3125 mol

Slide 42

2H _{2(g)}	+ O _{2(g)}	2 H ₂ O _(g)
I 1.0 mol	0.3125 mol	0 mol
C -2(0.3125)	-0.3125	+2(0.3125)
E 0.375	0	0.625 mol

Total moles gas= 0.375 + 0.625 = 1 mol

Slide 43

$$PV = nRT$$
$$P(10\text{ L}) = 1\text{ mol} \cdot \frac{0.082056\text{ L}\cdot\text{atm}}{\text{mol}\cdot\text{K}} \cdot 823.15\text{ K}$$
$$P = 6.8\text{ atm}$$
